PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6:

C09K 3/00

A1

(11) International Publication Number: WO 99/40162

(43) International Publication Date: 12 August 1999 (12.08.99)

US

(21) International Application Number: PCT/US99/02621

(22) International Filing Date: 8 February 1999 (08.02.99)

(30) Priority Data: 09/021,000 9 February 1998 (09.02.98)

(71) Applicant: CHEM LAB PRODUCTS, INC. [US/US]; Suite 1310, 300 East Second Street, Reno, NV 89501 (US).

(72) Inventors: KINCAID, Thomas, R.; 4850 Ramcreek Trail, Reno, NV 89509 (US). SOMESLA, Dana, W.; 3578 Mount Vernon Avenue, Riverside, CA 92502 (US).

(74) Agents: PALMER, John et al.; Ladas & Parry, Suite 2100, 5670 Wilshire Boulevard, Los Angeles, CA 90036 (US). (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published

With international search report.

(54) Title: A QUICKLY DISSOLVING CHLORINE RELEASE MIXTURE

(57) Abstract

A quickly dissolving chlorine release mixture for shock treatment of a substantial body of water as the consequence of solution in a localized region of a said body of water and subsequent dispersion into its entire volume. The mixture comprises: dry trichloroisocyanuric acid ("trichloro") particles having an exposed surface; and dry water soluble surfactant particles, said trichloro and surfactant particles being intimately and loosely mixed with one another. The surfactant has the properties of ready solution into the water and wetting of the surface of the trichloro particles, the surfactant being provided in an amount sufficient to wet the surfaces while the particles are in the localized region, and insufficient in amount to adversely affect the water when dispersed into its entire volume. The surfactant may comprise sodium lauryl sarcosinate or potassium perfluoroalkyl sulfonate, or their mixture.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	ТJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	iL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JР	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	zw	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	ΚZ	Kazakstan	RO	Romania		
cz	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	•	LI	Liechtenstein	SD	Sudan		
	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia	LK	LIUCIIA	30	o8.40.0		
1							

A QUICKLY DISSOLVING CHLORINE RELEASE MIXTURE

Field of the Invention

This invention relates to the "shock" treatment of substantial bodies of water with chlorine quickly released from trichloroisocyanuric acid. Trichloroisocyanuric acid is ordinarily very slow to dissolve and to release its chlorine.

Background of the Invention

Substantial bodies of water such as swimming pools, spas, and cooling towers require constant attention to the elimination of undesirable organic material such as waste material, organic compounds, algae and bacteria. The infusion of gaseous chlorine or of hypochlorite ion is a conventional way to provide an oxidizing agent for this purpose.

There are two levels of attack. One is for the long term maintenance of an existing agreeable condition, in which the level of the oxidizing agent is kept low, and the substance is supplied at a slow steady rate. Examples are the use of slowly dissolving tablets of the oxidizing agent placed in the body of water or in a recirculating system supplying it, or in a slow but steady stream of an oxidizing substance, such as gaseous chlorine, hypochlorite solution, oxygen, peroxides and ozone.

The other level of attack is when the conditions in the body

of water have deteriorated to the point where a slow supply is insufficient to eliminate the problem, and in which a vigorous attack must be made. In such event, a large amount of oxidizing agent is added to the body very quickly, so that the chlorine concentration rises well above that which would be maintained during routine operations. Such a treatment is called a "shock" treatment by persons familiar with this art.

Especially for domestic bodies of water, but also for many other substantial-sized bodies of water such as cooling towers, the amount of chemicals needed for a shock treatment is relatively small in weight and bulk, and would most conveniently be provided in small containers or pouches. These must be packaged to have a reasonable shelf life, and be agreeable when the package is opened. The best situation would be if the product were safe to store and handle as well as easy to use.

There are many products which from a chemistry standpoint can be used for shock treatment. Aqueous hypochlorite solutions are a familiar example. A pervasive problem is the bulk which must be handled, and the rapid loss of chlorine. Sources of treatment gas such as chlorine and ozone require capital installations and handling of hazardous materials.

In contrast, trichloro compounds would be preferable because of their substantial chlorine content, safety in handling and convenience. Still they are slow to dissolve.

This property of slow dissolving is ideal for long-term maintenance, and is the principal reason why trichloroisocyanuric acid (hereinafter called "trichloro") is so frequently used in compressed tablets kept submerged in a swimming pool or in a feeder. The trichloro in a tablet form will dissolve very slowly, and will maintain a steady, slow chlorine release for a long period of time when that characteristic is desirable. However, its slow dissolving rate, even in finely divided form, is a contraindication against its use for shock treatment because immediate chlorine release is the objective. An additional complication occurs because trichloro when finely grained simply floats and coagulates on the surface of the water for a long period of time, and when finally wetted, the clusters sinks to the bottom where they dissolves slowly.

Trichloro has a substantial advantage in its stability while dry, and in its low flammability. Except for the fact that unassisted it is useless for a shock compound, it would be an ideal domestic source of chlorine for this application.

In contrast, dichloroisocyanurate (hereinafter called "dichloro") is readily soluble, and is used for shock treatments. Its disadvantages compared to other chlorine compounds include, but are not limited to a higher price, lower relative available chlorine and the potential fire hazardous of the anhydrous form. Its only comparative advantage is its speed of solution into

water, which is rapid. For this reason it is used for shock treatments despite its disadvantages.

This situation has not gone unnoticed, and efforts have been made to provide mixtures of trichloro and other chemicals, which provide pathways by which the chlorine in trichloro can quickly be released. These arrangements have brought with them a number of problems, because characteristically they include substances which will react with trichloro to reduce it to the more soluble dichloro when in the water, or which provide other reaction paths involving reactants that must be stored together with the trichloro so as to be available for the intended reaction.

Accordingly, when used, trichloro has generally been accompanied by substances which bring their own expense and problems. The result of all of this is that trichloro has generally been disfavored for use in shock treatment of large bodies of water.

It is an object of this invention to provide a mixture of trichloro and a small amount of another substance to be described which mixture is stable, safe to handle, does not constitute a flammability risk, has a relatively lower toxic risk and which dissolves very quickly.

This composition is intended for use in substantial volumes of water in which the composition will be dissolved in an unbounded but very local region, from which it disperses into the larger volume. It is not intended as a local oxidizer or

cleanser for surfaces, or as a solution for direct use on anything, but instead is intended for infusion into the substantial volume very quickly after the trichloro is dissolved, thereby to provide a quick increase in chlorine in the substantial body of water.

There is yet another problem inherent in the use of trichloro. It has so much chlorine content per unit of weight or volume that a small amount of it suffices for a very large body of water. For example, 6 ounces of trichloro will suffice to shock treat a swimming pool containing 12,000 gallons of water, raising its chlorine concentration by about 3 ppm.

The small size of this portion presents the problem of providing to the consumer an amount of the product which can be used at one (1) pound per 12,000 gallons of water (a traditional premium consumer product dosage rate) for shock treatment.

Accordingly, a bulking agent may be provided as an inert diluent so a dose can be presented to the consumer in a single package and in a condition such that he can readily transfer it from the package, such as a plastic pouch, to the body of water. In addition the dose delivered is accurate, much more accurate than if a small amount of the trichloro were measured out of a larger container.

The bulking agent is inert relative to the dissolving of the trichloro into the water, and to its release of oxidizing agent.

Its purpose is not to assist these functions, but to dilute the trichloro by weight for both dose control and affixing an acceptable price per pound. However if this inert diluent can also provide other advantages such as reduction (or practical elimination) of flammability or to stabilize the active ingredients, or clarify the water (or condition the water) so much the better.

It is an object of this invention to accompany trichloro with a substance which will accelerate its solution without taking part in the chlorine release, and if desired, to provide additives as bulking agents, and other additives which provide other water treatment properties in addition to those provided by the released chlorine, but which take no part in the release of the chlorine or dissolving of the trichloro.

Brief Description of the Invention

This invention comprises a dry mixture of granulated trichloroiscyanuric acid ("trichloro") and a surfactant having the property when placed in water of wetting the trichloro and faciliting its quick solution into the water.

It is essential to this invention that the solution of trichloro be very rapid, and be completed during the time when the local concentration of the surfactant remains high enough adequately to disperse and wet the surface of trichloro granules when it is exposed to water during its immersion and dissolution.

The same amount of surfactant dispersed through the entire body of water would not suffice for the purpose. The trichloro would simply stay dry or dissolve very slowly as though the surfactant were not present at all. To provide surfactant in the whole body of water in a concentration sufficient to dissolve the product would be to make the water system foamy and unacceptable for its intended purpose. This is a local action in a limited but unbounded region of a much larger body of water.

According to this invention, the surfactant and the trichloro are dry loose particles that are intimately and physically mixed, but not otherwise pressed or tabletted. A very large exposed area of the trichloro is desired, a result attained by making the particle size small. The mixture is dumped together onto the surface of the water. The surfactant dissolves almost instantly to form a small local unbounded region containing a sufficiently high concentration of surfactant, within which the trichloro is present to dissolve the trichloro. The trichloro is wetted because of the surfactant, and quickly dissolves in the local region. The surfactant remains at an adequate concentration in this region so that as the trichloro particles disperse, their newly exposed surfaces are wetted and dissolved. After dissolution is completed, the surfactant and the dissolved trichloro disperse into the larger body, where the classical release of chlorine occurs. The concentration of

surfactant in the body of water is very small in the total body of water, on the order of a 100,000 fold dilution, while the chlorine concentration is increased to an effective level for shock treatment.

It is instructive to observe this event. A packet of this mixture dumped onto the surface of a swimming pool swiftly forms a downwardly and laterally progressing enlarging plume in which the presence of the trichloro as it dissolves and disperses is visually evident. This is the "region". The plume soon disperses, and there is no residual solid trichloro. In a short time, usually much less than a minute, all of the trichloro will have dissolved. There is no visible indication that anything had been added.

Detailed Description of the Invention

The source of available chlorine in the mixture according to this invention is trichloro-s-triazine trione, also called trichloroisocyanuric acid ("trichloro" herein). It is the fully chlorinated product obtained from the tri-sodium salt of isocyanuric acid. When dissolved, it reacts with water and releases available chlorine used for oxidizing purposes.

As discussed above, its unassisted rate of solution in water is unacceptably slow for use in shock treatment. The inventors herein have observed that the rate of solution will be accelerated by providing conditions in which a sufficient

concentration of a suitable surfactant wets the trichloro, and will keep the surface of the trichloro wet while it dissolves.

Obviously this cannot be a pre-wet combination, because when the trichloro is wetted, it immediately begins its decomposition.

There have been described arrangements in which chemical pathways are provided to accelerate the dissolution of trichloro in water. These require the use of a large percentage of a compound which is reactive with the trichloro. This poses the problems of cost, and maintaining the product while in storage in a suitable dry condition.

In order to be available for use when the product is place in water, it is evident that the surfactant must not react with the trichloro in the packet. Such surfactants appear to be very few in number. For this product, sodium lauroyl sarcosinate is suitable and preferred. It is very soluble, and provided in powder form. Potassium perfluoroalkyl sulfonate may be used instead, or in combination with it to provide the same relative amount by weight.

In order to assure that there will be a sufficient local and temporary concentration of surfactant to be present while the trichloro is dissolving, the ratio by weight of dry surfactant to dry trichloro should be between about 1300:1 and about 130:1, preferably about 600:1.

Both must be dry and are best present as a mixture of

granules or particles so as to minimize any tendency to float on the surface of the water, as might occur if they are too finely powdered, or to sink before dissolving if too large a particle.

For the trichloro, a particle size to pass 20 mesh and 50% to be retained on 200 mesh screen is suitable. The particle size of the surfactant may be about the same, but is less critical because it dissolves so quickly.

This invention is a binary mixture which consists essentially of a dry mixture of trichloro and a suitable non-reactive surfactant. Inert additives, which are inert in the sense that they take no part in the release of chlorine from the trichloro may be added for various other purposes.

The principal use of such additives is as a bulking agent, where a small enough amount of the active ingredients (trichloro and surfactant) can be presented to the consumer in a larger bulk at a lower cost per pound with significantly reduced chemical hazard. These additives must not themselves be deleterious to the mixture in storage or in the water, nor to the quality of the water. For example they should not cause cloudiness, or leave a residue in the water or on the bottom.

For the above reasons, the additives are preferably soluble and must be stable relative to the rest of the mixture. In a commercially viable product, they should also be inexpensive.

One useful additive for a bulking agent is sodium chloride,

which meets all requirements, except that it is slightly hygroscopic, so that great care must be taken during packaging. When it is used, it is better to accompany it with a substance which acts as a desiccant which will preferentially absorb any water, and which can absorb any slight chlorine odor which maybe generated in the package. For this purpose, sodium carbonate is very suitable. It acts as a desiccant and as an odor-collector.

While by its very presence boric acid acts as a bulky agent, it is included in this formulation for two others of its well-known properties. The first is as a flame retardant, an advantage during handling and storage. The other is to improve flow ability of the mixture. This assists in assuring complete mixing, accurate packaging, and ready release from the package.

The amount of surfactant used in this invention is insufficient for any practical use except within the limited region in which it is placed as part of its mixture with the trichloro, and is deliberately made so small that after dispersion its concentration is so trifling that it will not grossly affect the pool conditions.

The mixture according to this invention consists essentially of trichloro and surfactant in the ratio between themselves about 1300 to 130 parts by weight of trichloro to about 1 part by weight of surfactant. The preferred amount of surfactant is that which will provide the minimum concentration

of surfactant in the limited region that will result in quick solution of the trichloro. While amounts somewhat greater than this will do no harm to users of the treated water, still it is best to limit the amount of a material which in a sufficient concentration could produce foaming in larger concentrations.

The presently preferred formulation, of which 6 ounces will provide about 3 ppm of chlorine in a 12,000 gallon pool, is as follows (in percentages by weight):

Boric acid, dry powder: about 10%

Sodium carbonate, dry powder: about 10%

Sodium chloride, dry powder: about 40%

Surfactant, sodium lauroyl-

sarcosinate dry powder: about 0.015%

Trichloro: balance to 100%, in

which all except the trichloro and surfactant are bulking agents. While only one or two of these bulking agents can be used, the preferred formulation will use all three. Boric acid has the additional optional properties of improving flowability and flame retardancy.

The ratio of bulking agent or agents may vary from this total of 60% by weight, thereby enabling a packet of the same weight to contain more or less of the trichloro. It has been found that if at least a total of about 20% by weight of the above mixture is of these bulking agents, there is a useful control of inadvertent

dampness, resulting odor, and some reduction in flammability. Flammability of trichloro is very low to begin with, so that boric acid is optional. There is no upper limit, except that there must be a sufficient amount of the trichloro to provide an advantage. The basic mixture of trichloro and surfactant is fully useful and safe, without bulking agents or additives for other purposes such as odor control, desiccation, or reduction of flammability (while also serving as bulking agents).

This invention is not to be limited by the embodiments described in the description, which are given by way of example and not of limitation, but only in accordance with the scope of the appended claims.

I CLAIM:

1. A quickly dissolving chlorine release mixture for shock treatment of a substantial body of water as the consequence of solution in a localized region of a said body of water and subsequent dispersion into its entire volume, said mixture consisting essentially of:

dry trichloroisocyanuric acid ("trichloro") particles having an exposed surface; and

dry water soluble surfactant particles, said trichloro and surfactant particles being intimately and loosely mixed with one another;

said surfactant having the properties of ready solution into the water and wetting of the surface of the trichloro particles, said surfactant being provided in an amount sufficient to wet said surfaces while said particles are in said localized region, and insufficient in amount to adversely affect the water when dispersed into its entire volume.

2. A mixture according to claim 1 in which said mixture further includes an inert water soluble bulking agent mixed into it whereby to provide a more conveniently marketed and dispensed product, said bulking agent being ineffective substantially to change the rate of solution of the trichloro, or to affect its release or rate of release of chlorine.

3. A mixture according to claim 1 in which the ratio by weight of said acid to surfactant is between about 130:1 and about 1300:1

- 4. A mixture according to claim 3 in which said ratio is about 600:1.
- 5. A mixture according to any one of claims 1-4 in which said surfactant is sodium lauroyl sarcosinate or potassium perfluoroalkyl sulfonate, or their mixture.
- 6. A mixture according to any one of claims 1-4 in which said trichloro particles pass a 20 mesh screen and about 50% are retained on a 200 mesh screen.
- 7. A mixture according to claim 2 in which said bulking agent comprises a desiccant.
- 8. A mixture according to claim 2 in which said bulking agent comprises an odor control for absorbing chlorine odor.
- 9. A mixture according to claim 2 in which said substance to improve flowability is included in said bulking agent.

10. A mixture according to claim 9 in which said substance is boric acid.

11. A quickly dissolving chlorine release mixture for shock treatment of a substantial body of water, consisting essentially of, in percentages by weight:

dry trichloroisocyanuric acid
 ("trichloro") particles:

balance to 100%

dry sodium lauroyl sarcosinate:

about 0.015%

and, bulking agents ineffective substantially to change the rate of solution of the trichloro or to affect its release or rate of release of chlorine:

about 60%.

12. A mixture according to claim 11 in which said bulking agents are the following, in the percentages by weight indicated:

Boric acid, dry powder:

about 10%

Sodium carbonate, dry powder:

about 10%

Sodium chloride, dry powder

about 40%.

13 A mixture according to claim 11 in which said trichloro particles pass a 20 mesh screen and about 50% are retained on a 200 mesh screen.

15. A mixture comprising, in percentages by weight: trichloroisocyanuric acid

("trichloro"):

balance to 100%;

sodium lauroyl sarcosinate:

about 0.015%;

and, a bulking agent:

about 20 to about 60%.

- 16. A mixture comprising about 20% to about 60% by weight of a bulking agent, the balance of the mixture to 100% further comprising trichloroisocyanuric acid ("trichloro") and sodium lauroyl sarcosinate, wherein the ratio by weight of the acid to the sodium lauroyl sarcosinate ranges from about 1300:1 to about 130:1.
- 17. A mixture as claimed in claim 2, wherein the bulking agent is about 20% to about 60% by weight of the mixture, wherein the balance of the mixture to 100% consists essentially of the acid and the surfactant, and wherein the ratio by weight of the acid to the surfactant ranges from about 1300:1 to about 130:1.

INTERNATIONAL SEARCH REPORT

International application No. PCT/US99/02621

		<u></u>						
A. CLASSIFICATION OF SUBJECT MATTER IPC(6) : CO9K 3/00								
US CL: 252/190 According to International Patent Classification (IPC) or to both national classification and IPC								
		national classification and IPC						
	DS SEARCHED ocumentation searched (classification system followed)	hy classification symbols)						
		by classification symbols)						
U.S. :	252/190, 187.1, 187.33, 187.34							
Documentati	ion searched other than minimum documentation to the	extent that such documents are included	in the fields searched					
	lata base consulted during the international search (na	me of data base and, where practicable	e, search terms used)					
C. DOCUMENTS CONSIDERED TO BE RELEVANT								
Category*	Citation of document, with indication, where ap	propriate, of the relevant passages	Relevant to claim No.					
Y	US 5,128,342 A (GLOBUS) 07 July 19	1, 2, 5-8, 11, 13						
Y	US 4,557,926 A (NELSON et al) 10 December 1985, col. 1, lines 1,2,5 60-68; col. 2, lines 1-32.							
A	US 4,193,888 A (MCHUGH) 18 Marc							
A	US 5,114,647 A (LEVESQUE et al) 1							
A	US 5,055,219 A (SMITH) 08 October							
A	US 4,279,764 A (BRUBAKER) 21 Jul							
Α	US 4,863,632 A (ARONSON et al) 05 September 1989.							
	ther documents are listed in the continuation of Box C							
.V. q	pecial categories of cited documents: ocument defining the general state of the art which is not considered b be of particular relevance	"T" later document published after the int date and not in conflict with the app the principle or theory underlying th	lication but cited to understand					
E 0	arlier document published on or after the international filing date locument which may throw doubts on priority claim(s) or which is	"X" document of particular relevance; the considered novel or cannot be considered when the document is taken alone	e claimed invention cannot be ared to involve an inventive step					
C:	ited to establish the publication date of another citation or other pecial reason (as specified) locument referring to an oral disclosure, use, exhibition or other	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination						
P d	neams locument published prior to the international filing data but later than	being obvious to a person skilled in the art *A* document member of the same patent family						
	he priority date claimed e actual completion of the international scarch	Date of mailing of the international se	earch report					
	RCH 1999	15 APR 1999	\bigcirc					
		Authorized officer Share How How						